

ORIGINAL

## The treatment with podiatric massage for individuals with plantar fasciitis

### El tratamiento con masaje podálico a personas con fascitis plantar

Odalys Senado Pedrosa<sup>1</sup>  

<sup>1</sup>Centro de Investigaciones Clínica. Departamento de Fisioterapia y Rehabilitación. La Habana. Cuba.

Cite as: Senado Pedrosa O. The treatment with podiatric massage for individuals with plantar fasciitis. Interdisciplinary Rehabilitation / Rehabilitación Interdisciplinaria. 2026; 6:222. <https://doi.org/10.56294/ri2026222>

Submitted: 28-04-2025

Revised: 20-08-2025

Accepted: 30-12-2025

Published: 01-01-2026

Editor: PhD. Nicola Luigi Bragazzi 

Corresponding author: Odalys Senado Pedrosa 

#### ABSTRACT

**Introduction:** in the treatment of plantar fasciitis, foot massage is effective due to its variety of manipulations, gentle movements on the soft tissues of the body, by acting with deep and energetic pressure. These maneuvers are essential and adaptable according to the individual needs of each person with the aim of providing relief and well-being.

**Objective:** to diagnose the initial state for the execution of foot massage to people with plantar fasciitis at the Dr. Cosme Ordoñez Carceller University Polyclinic.

**Method:** a descriptive, cross-sectional study was carried out, theoretical methods were applied, the systematization of references related to the subject and empirical methods such as a survey of people with plantar fasciitis, to identify the type of foot massage treatment applied, physical examination and diagnostic tests to determine their level of affectation.

**Results:** analysis of the results of the instruments used to assess the initial diagnosis and their methodological triangulation allowed us to identify problems and potential opportunities for the application of foot massage in patients with plantar fasciitis attending this clinic.

**Conclusions:** the problems and potential opportunities identified allowed for the development of the protocol.

**Keywords:** Plantar Fasciitis; Foot Massage; Podiatry.

#### RESUMEN

**Introducción:** en el tratamiento de la fascitis plantar, el masaje podálico es efectivo por su variedad de manipulaciones, movimientos suaves en los tejidos blandos del cuerpo, al actuar con presiones profundas y energéticas. Estas maniobras son esenciales y adaptables según las necesidades individuales de cada persona con el objetivo de brindar alivio y bienestar.

**Objetivo:** diagnosticar el estado inicial para la ejecución del masaje podálico a personas con fascitis plantar en el Policlínico Universitario Dr. Cosme Ordoñez Carceller.

**Método:** se efectuó un estudio, descriptivo, transversal, se aplicaron métodos teóricos la sistematización de referentes relacionados con el tema y métodos empíricos como encuesta a personas con fascitis plantar, para identificar el tipo del tratamiento con masaje podálico que se aplica, la exploración física y pruebas diagnósticas para determinar su nivel de afectación.

**Resultados:** el análisis de los resultados de los instrumentos que se aplicaron para la valoración del diagnóstico inicial y su triangulación metodológica, permitieron identificar problemas y potencialidades para la aplicación del masaje podálico en personas con fascitis plantar, quien asisten a este policlínico.

**Conclusiones:** los problemas y potencialidades que se identificaron permitieron la estructuración del protocolo.

**Palabras clave:** Fascitis Plantar; Masaje Podálico; Podología.

## INTRODUCTION

Plantar fasciitis (PF) is inflammation of the plantar fascia. The pain is localized to the inner bottom of the heel and can extend to the toes.<sup>(1,2)</sup> PF is the most common condition treated by podiatrists and the most common cause of calcaneal pain. It is estimated to affect 2 million Americans annually and results in more than 1 million clinic visits,<sup>(3)</sup> where treatment with foot massage is used.

Massage as a therapeutic technique is used to relieve pain and improve health. In the treatment of PF, massage is effective in reducing symptoms. This method focuses on applying intense and sustained pressure to the affected area. Frictional movements along the plantar fascia release accumulated tension and reduce muscle contractures in the sole of the foot and surrounding tissues.<sup>(4)</sup> Musculoskeletal disorders affect the locomotor system and can manifest acutely and transiently, presenting with pain and limited mobility. These conditions can involve muscles, tendons, and bones, negatively impacting people's well-being and ability to participate in society,<sup>(5)</sup> including PF.

In 2023, the University Polyclinic Dr. Cosme Ordoñez Carceller, with a population of approximately 16,490 inhabitants, saw an increase in the number of people with PF attending Podiatry services. It is estimated that the age of debut (40-60 years) was the highest, with the female sex representing the highest incidence.

This background made it possible to identify the scientific problem of determining the need for rehabilitative treatment with foot massage for people with PF and the research objective: to diagnose the initial state with foot massage for people with PF.

## METHOD

A descriptive, cross-sectional study was conducted in podiatry services between March and May 2024.

### Inclusion criteria

Selection of people with PF attending podiatry services.

### Exclusion criteria

Diseases where foot massage is contraindicated.

### Diagnosis of the initial state

#### *Parameterisation of the variable*

The information was collected from a population of 15 people (100 %) who attended Podiatry services between September 2023 and diagnosed with PF in the characterization of the sample. The sample included six men (40 %) and nine women (60 %) aged 40-60.

The parameterization procedure for executing the initial diagnostic stage is explained, followed by analyzing the results of each instrument applied to arrive at the inventory of problems and potential.<sup>(6,7,8)</sup> A single variable, with two dimensions and 23 indicators, is identified: foot massage for people with VF.

*Diagnostic dimension:* stage in which the symptoms and signs of people with PF are assessed.

- Pain threshold level at rest
- Pain threshold level on palpation
- Level of pain threshold level on the footrest
- Level of stiffness (joint mobility)
- Level of swelling in the affected area
- Level of numbness in legs and feet
- Level of difficulty in maintaining correct posture
- Level of stiffness in the arch of the foot
- Level of difficulty in walking
- Level of pain in the sole
- Level of pain in the heel
- Level of stiffness in the Achilles tendon
- Level of stiffness in the gastrocnemius muscles
- Level of relief when applying foot massage

*Procedural-rehabilitative dimension:* stage of rehabilitative treatment where foot massage is applied to people with PF.

- Level of execution of strength exercises
- Level of execution joint mobility
- Level of execution of friction
- Level of execution of rubbing
- Level of execution of kneading
- Level of execution of percussion
- Execution level stretching
- Level of execution shaking
- Level of reflexology execution

Theoretical methods, such as documentary analysis and systematization, and empirical methods, such as surveys of people with PF, observation of treatment with foot massage, physical examination, diagnostic tests of PF (pain and flexibility), the Numerical Verbal Pain Scale (ENVD), and the Simple Verbal Pain Scale (EVSD) were applied.

#### *Statistical methods*

Descriptive statistics were used to calculate the percentages in the procedure for the data obtained in the empirical investigations.

#### *Methodological triangulation technique*

To process and integrate the results obtained from the inventory of problems and potentialities.

## RESULTS

Assessment of the survey of people with VET:

| Table 1. Decision                                                                                            |              |
|--------------------------------------------------------------------------------------------------------------|--------------|
| <b>Decision criteria (indicators 1 to 13)</b>                                                                |              |
| If more than 60 % of the collected data is found in the sum of the norms YES and FREQUENTLY, it is assessed: | Problem      |
| If the data collected is less than 59 % in the sum of the normotypes YES and FREQUENTLY it is assessed:      | Satisfactory |
| <b>Decision criteria (indicators 14 to 17)</b>                                                               |              |
| When the data collected is more than 60 % in the sum of the normotypes YES and FREQUENTLY it is assessed:    | Satisfactory |
| If the data collected is less than 59 % in the sum of the normotypes YES and FREQUENTLY it is assessed:      | Problem      |

**Table 2.** Assessment of the survey results

| No. | Diagnostic Criteria                             | Yes         | Sometimes   | No         |
|-----|-------------------------------------------------|-------------|-------------|------------|
| 1   | Painful at rest (1.1)                           | 5(33,33 %)  | 7(46,67 %)  | 3(20 %)    |
| 2   | Pain on palpation (1.2)                         | 12(80,00 %) | 3(20,00 %)  | -          |
| 3   | Pain on foot rest (1.3)                         | 12(80,00 %) | 3(20,00 %)  | -          |
| 4   | Stiffness in the foot (1.4)                     | 10(66,67 %) | 3(20,00 %)  | 2(13,33 %) |
| 5   | Swelling in the affected area (1.5)             | 13(86,67 %) | 2(13,33 %)  | -          |
| 6   | Numbness in the legs and feet (1.6)             | 7(46,67 %)  | 7(46,67 %)  | 1(6,67 %)  |
| 7   | Difficulty in maintaining correct posture (1.7) | -           | 10(66,67 %) | 5(33,33 %) |
| 8   | Stiffness in the arch of the foot (1.8)         | 9(60,00 %)  | 5(33,33 %)  | 1(6,67 %)  |
| 9   | Difficulty walking (1.9)                        | 10(66,67 %) | 5(33,33 %)  | -          |
| 10  | Feeling plantar pain (1.10)                     | 8(53,33 %)  | 7(46,67 %)  | -          |
| 11  | Pain in the heel (1.11)                         | 7(46,67 %)  | 8(53,33 %)  | -          |
| 12  | Achilles tendon tightness (1.12)                | 8(53,33 %)  | 6(40,00 %)  | 1(6,67 %)  |
| 13  | Stiffness in the gastrocnemius muscle (1.13)    | 8(53,33 %)  | 6(40,00 %)  | 1(6,67 %)  |
| 14  | Feels relief with foot massage (1.14)           | 13(86,67 %) | 2(13,33 %)  | -          |

The most affected criteria are those related to pain (1.1,1.2,1.3,1.10,1.11); swelling (1.5) and stiffness (1.4,1.8,1.12,1.13). The criteria related to difficulty in maintaining balance, correct posture and walking (1.7,1.9) are due to stiffness and pain. However, most of the time they feel relief with massage (1.14).

**Table 3.** Participant observation of podalic massage treatment for people with VF

| No. | Criteria according to observation            | SO         | SAV         | NSO         |
|-----|----------------------------------------------|------------|-------------|-------------|
| 1   | Strength exercises are performed (2.1)       | 5(33,33 %) | -           | 10(66,67 %) |
| 2   | Joint mobility exercises are performed (2.2) | 5(33,33 %) | 8(66,67 %)  | 2(13,33 %)  |
| 3   | Surface Friction is performed (2.3)          | 5(33,33 %) | 10(66,67 %) | -           |
| 4   | Rubbing is performed (2.4)                   | 5(33,33 %) | 10(66,67 %) | -           |
| 5   | Kneading is performed (2.5)                  | 5(33,33 %) | 10(66,67 %) | -           |
| 6   | Percussion is performed (2.6)                | 5(33,33 %) | -           | 10(66,67 %) |
| 7   | Stretching is executed (2,7)                 | 5(33,33 %) | 2(13,33 %)  | 8(53,33 %)  |
| 8   | Shaking is performed (2,8)                   | 5(33,33 %) | 2(13,33 %)  | 8(53,33 %)  |
| 9   | Reflexology is performed (2,9)               | 3(20,00 %) | -           | 12(13,33 %) |

**Table 4.** Participant observation of podalic massage treatment for people with PF with percentage

| Decision criteria                                                                                                   | Ranking           |
|---------------------------------------------------------------------------------------------------------------------|-------------------|
| When the data collected is less than 60 % in the SO normotype it is assessed:                                       | Problem           |
| When the data collected is between 61 % and 69 %. in the SO normotype is assessed:                                  | In transformation |
| When the maximum expected maximum of the collected data is between 70 % and 100 %. in the SO normotype is assessed: | Satisfactory      |

Foot reflexology massage is generally applied, but a methodology is needed to improve the application of this treatment. Foot reflexology needs to be applied more to improve the health of people suffering from PF. The most used manipulations are stretching, rubbing, and kneading, and the least used is percussion.

Strength and joint mobility exercises are carried out, to which other exercises should be added.

Table 5. Diagnostic test for PF (pain and flexibility)

| Sitting Position     | Movement                                                      | Valuation                                                      | Anatomical region involved | Positive result (bad) | Negative result (good) |
|----------------------|---------------------------------------------------------------|----------------------------------------------------------------|----------------------------|-----------------------|------------------------|
| Knee extended        | Dorsal flexion of the ankle<br>(1,4,1,6,1,10,1,11, 1,12,1,13) | Normal range of motion 10°. positive with restriction and pain | Triceps suralis            | 15                    | -                      |
| With knee flexed 90° | Dorsal flexion of the ankle<br>(1,4,1,6,1,10,1,11, 1,12,1,13) | Normal range of motion 10°. positive restriction and pain      | Complex (gastro-oil)       | 15                    | -                      |

### Description of the maneuver

In the supine position with the knee flexed or extended. In both cases, the examination is performed on the hindfoot joints in a reduced position to block them and avoid measurement errors.

### With the knee extended

Assessment of the dorsal flexion of the ankle and the involvement of the triceps suralis, normal range of motion of dorsal flexion 10°. It is repeated with the knee flexed to 90°; if there is retraction or contracture with flexion of the knee, the limitation of dorsal flexion remains; the retraction would be due to the soleus muscle. In both cases, the test is considered positive as it would present a restriction of the gastro-soleus complex.

### Position of the person

Seated with hip flexion at 90°. The ankle is stabilized with the passive hand while, with the active hand, the proximal phalanx of the first toe (hallux) is grasped, performing dorsal flexion. This movement produces flexion of the first metatarsal and an increase in the plantar arch, causing tension in the central band of the plantar fascia. We then palpate along the fascia. The test is positive if it reproduces the pain.

*Positive:* there is a restriction of movement and pain.

*Negative:* normal movement and no pain

All the people who had this diagnosis were assessed as positive.

### Numerical Verbal Pain Scale (ENVD)

Scores from (0 to 10) your degree of pain, and is divided into four groups: 0 no pain, (1 to 3) mild pain, (4 to 6) moderate pain, (7 to 10) very severe pain.




Figure 1. Numerical Verbal Pain Scale (ENVD)

### Pain is assessed as

No pain 0, mild pain 2, moderate pain 5, and severe pain 8.

### Physical examination of people receiving foot massage treatment with PF

Is done to determine the existence or absence of physical alterations, as well as an analysis of the signs and symptoms of the people produced by the disease.

The physical examination results aimed at people with PF showed a distribution of forefoot 1 (6,67 %), midfoot 4 (26,67 %), rearfoot 8 (53,33 %), and malleolus 2 (13,33 %).

More than 50 % of the physical alterations were identified in the rearfoot region, which coincides with the studied references.

### Results of the methodological triangulation

Methodological triangulation makes it possible to group and compare the information received from different sources, techniques, and instruments to identify coincidences and discrepancies in the phenomenon under study.<sup>(9)</sup>

When the results of the instruments applied are compared, it is observed that the indicators related to pain, as well as those related to joint stiffness and mobility, are valued as problems; therefore, dimension two (diagnosis) is valued as a problem.

The hindfoot region (calcaneus) is the most affected. Few manipulations of podsollic massage, strengthening exercises, and joint mobility are appreciated; people with PF accept the message, so it is necessary to improve this technique. Some indicators of dimension two, rehabilitative procedural dimension 2, are difficult, but they are valued as part of the transformation process.

## **DISCUSSION**

Podiatry treats pathologies that manifest themselves with painful symptomatology, establishing a complementary relationship to reduce foot discomfort, which is essential for identifying functional and structural alterations. This requires a thorough assessment and an individualized examination to optimize the health of individuals.<sup>(10)</sup>

Musculoskeletal disorders are diseases characterized by a pathology involving injuries to bones, muscles, tendons, nerves, joints, or ligaments. These conditions can be very painful and cause chronic problems with motor functions.<sup>(11)</sup> Neuromusculoskeletal disorders in the feet encompass a set of alterations of the locomotor apparatus of pathological origin, which include foot problems such as aches and pains.

The types of massage include therapeutic massage, which improves circulation, restores mobility, and reduces ailments; relaxing massages, which provide comfort and regulate tension; and preventive massages, which relax tense areas and prevent injuries.<sup>(12)</sup>

## **CONCLUSIONS**

Parameterisation of the object of study was carried out to apply the initial diagnosis related to the treatment with foot massage aimed at people with PF. Problems and potentialities were identified and considered for the elaboration of the action protocol.

## **ACKNOWLEDGEMENTS**

Dr.C. Tamara Rodríguez Morera, Dr.C. Rafael Rousseau Yera.

## **REFERENCES**

1. Servicios A. ¿Qué es la talalgia y cómo curarla en mayores? [Internet]. Aiudo Servicios. 2024 [citado 5 de mayo de 2025]. Disponible en: <http://servicios.aiudo.es/podologia/que-es-talgia-tratamiento-diferencia-fascitis-plantar-ejercicios/>
2. Lagos M. FASCITIS PLANTAR: Información para pacientes y familiares. Hosp Clínico San Carlos [Internet]. noviembre de 2024 [citado 5 de mayo de 2025];(1):2. Disponible en: <https://www.comunidad.madrid/hospital/clinicossancarlos/sites/clinicossancarlos/files/inline-files/D30%20FASCITIS%20PLANTARV1%20NOV%202024.pdf>
3. Monara D, B. Parks C. McGraw Hill Medical. 2024 [citado 5 de mayo de 2025]. Fascitis plantar. Disponible en: <https://accessmedicina.mhmedical.com/content.aspx?sectionid=286823098&bookid=3463>
4. bigpiespro. Fascitis Plantar y Masajes: Beneficios y Técnicas para Aliviar la Tensión [Internet]. Podología en Cali | Big Pies. 2024 [citado 23 de abril de 2025]. Disponible en: <https://big-pies.com/fascitis-plantar-y-masajes-beneficios-y-tecnicas-para-aliviar-la-tension/>
5. Musculoskeletal health [Internet]. 2021 [citado 5 de mayo de 2025]. Disponible en: <https://www.who.int/news-room/fact-sheets/detail/musculoskeletal-conditions>
6. Freire DCEEE. Las variables y su operacionalización en la investigación educativa. Parte I. Rev Conrado [Internet]. 8 de octubre de 2018 [citado 18 de abril de 2025];14(65):39-49. Disponible en: <https://conrado.ucf.edu.cu/index.php/conrado/article/view/814>
7. Añorga Morales DrCJ, Valcárcel Izquierdo DrCN, Che Soler DrCJ. La parametrización en la investigación educativa. 45 Aniversario ISP “Enrique José Varona” [Internet]. diciembre de 2008 [citado 18 de abril de 2025];(47):25-32. Disponible en: <https://www.redalyc.org/articulo.oa?id=360635567005>
8. SOLIS SOLIS S. Modelo de evaluación del desempeño profesional del licenciado en higiene y epidemiología. 2017 [citado 6 de mayo de 2025]. Repositorio de Tesis en Ciencias Biomédicas y de la Salud de Cuba - Modelo de evaluación del desempeño profesional del licenciado en higiene y epidemiología. Disponible en: <https://>

tesis.sld.cu/index.php?P=FullRecord&ID=826

9. Vera C, Villalón CM. La Triangulación entre Métodos Cuantitativos y Cualitativos en el Proceso de Investigación TRIANGULATION BETWEEN QUALITATIVE AND QUANTITATIVE METHODS IN THE RESEARCH PROCESS Artículo de Educación. En 2005 [citado 10 de mayo de 2025]. Disponible en: <https://www.semanticscholar.org/paper/La-Triangulaci%C3%B3n-entre-M%C3%A9todos-Cuantitativos-y-en-Vera-Villa%C3%B3n/5cf49dc88c3c8b6ba4b19a7b25a2711f85cf1a22>

10. ClinicasOM. Relación entre fisioterapia y podología [Internet]. ClinicasOM. 2021 [citado 31 de marzo de 2025]. Disponible en: <https://clnicasom.com/fisioterapia-y-podologia-2/>

11. Rivera Guillén MA, Sanmiguel Salazar MF, Serrano Gallardo LB, Nava Hernández MP, Moran Martínez J, Figuerola Chaparro LC, et al. Factores asociados a lesiones músculo-esqueléticas por carga en trabajadores hospitalarios de la ciudad de Torreón, Coahuila, México. Cienc Amp Trab [Internet]. agosto de 2015 [citado 30 de marzo de 2025];17(53):144-9. Disponible en: [http://www.scielo.cl/scielo.php?script=sci\\_abstract&pid=S0718-24492015000200008&lng=es&nrm=iso&tlng=es](http://www.scielo.cl/scielo.php?script=sci_abstract&pid=S0718-24492015000200008&lng=es&nrm=iso&tlng=es)

12. Runner's World [Internet]. 2023 [citado 6 de marzo de 2025]. El masaje de 2 minutos que te ayudará a aliviar el dolor de la fascitis plantar. Disponible en: <https://www.runnersworld.com/es/salud-lesiones-runner/a2001394/fascitis-plantar-remedios/>

## FUNDING

None.

## CONFLICT OF INTEREST

None.

## AUTHORSHIP CONTRIBUTION

*Conceptualisation:* Odalys Senado Pedroso.

*Research:* Odalys Senado Pedroso.

*Methodology:* Odalys Senado Pedroso.

*Validation:* Odalys Senado Pedroso.

*Visualisation:* Odalys Senado Pedroso.

*Writing - original draft:* Odalys Senado Pedroso.

*Writing - revision and editing:* Odalys Senado Pedroso.